

Hilbert space inner products for PT-symmetric Su-Schrieffer-Heeger models

Frantisek Ruzicka *

*Faculty of nuclear sciences and physical engineering
Czech technical university in Prague
Brehova 7, 115 19 Prague
Czech Republic*

*Nuclear physics institute, Czech academy of sciences
Hlavni 130, 250 86 Husinec-Rez
Czech republic*

April 5, 2016

Abstract

A Su-Schrieffer-Heeger model with added PT-symmetric boundary term is studied in the framework of pseudo-hermitian quantum mechanics. For two carefully chosen special cases, a complete set of pseudometrics is constructed in closed form. When complemented with a condition of positivity, the pseudometrics determine all the physical inner products of the considered model.

1 Introduction

The Su-Schrieffer-Heeger (SSH) model [1, 2] is one of the benchmark topologically nontrivial models in physics of condensed matter. It provides a convenient description of certain physical systems [3, 4], and serves as one of the simplest examples of topological insulators [5]. The n -site SSH model may be expressed as

$$H_{SSH}^{(n)} = \sum_{i=1}^N \left[t(1 - \Delta \cos \theta) a_{2i-1}^\dagger a_{2i} + t(1 + \Delta \cos \theta) a_{2i}^\dagger a_{2i+1} + h.c. \right] \quad (1)$$

with $a_i^\dagger, a_i$ being the i -th site fermionic creation and annihilation operators. It has been suggested [2] to complement this hermitian SSH model with a non-hermitian $\mathcal{PT}$ -symmetric (invariant under simultaneous action of parity and time reversal) boundary term

$$H_I = \gamma a_1^\dagger a_1 + \gamma^* a_n^\dagger a_n \quad (2)$$

*Electronic address: ruzicfra@fjfi.cvut.cz

with $\gamma = i\alpha, \alpha \in \mathbb{R}$. Despite apparent non-hermicity of the resulting operator $H = H_{SSH} + H_I$, it has been shown that its eigenvalues remain real for $|\alpha|$ sufficiently small. This motivates further examination of this model from the viewpoint of $\mathcal{PT}$ -symmetric quantum mechanics.

$\mathcal{PT}$ -symmetric quantum mechanics [6, 7, 8] is a theoretical framework for studying representations of quantum observables by non-hermitian operators. It is based on a simple fact, that an observable may be represented by a non-hermitian operator on a given Hilbert space $\mathcal{H}^{(F)} = (\mathcal{V}, \langle \cdot | \cdot \rangle)$, as long as such operator is hermitian in another Hilbert space $\mathcal{H}^{(S)} = (\mathcal{V}, \langle \phi | \psi \rangle_\Theta)$ [9]. The inner products on $\mathcal{V}$ are in one-to-one correspondence with so-called metric operators Θ through the formula

$$\langle \phi | \psi \rangle_\Theta = \langle \phi | \Theta | \psi \rangle \quad (3)$$

where we require Θ to be bounded, hermitian and positive in order to generate a genuine inner product [10]. The hermicity condition of H in $\mathcal{H}^{(S)}$ could be expressed in operator form as

$$H^\dagger \Theta = \Theta H \quad (4)$$

which is a direct generalization of the standard condition $H = H^\dagger$. Operators H satisfying eq. 4 for at least one metric operator Θ are called quasi-hermitian [11]. It is clear that on a given Hilbert space $\mathcal{H}^{(F)}$, all admissible observables are described by quasi-hermitian instead of hermitian operators. Furthermore, recall that for any bounded, positive Θ , there always exists a decomposition $\Theta = \Omega^\dagger \Omega$ with Ω bounded. Inserting this decomposition into eq. 4 yields

$$h := \Omega H \Omega^{-1} = (\Omega H \Omega^{-1})^\dagger = h^\dagger \quad (5)$$

which shows that being quasi-hermitian is equivalent to being boundedly diagonalizable with real spectrum (instead of unitarily diagonalizable). However, the construction of the similarity transformation Ω may be in general very difficult for a given H . Special attention is usually given to parametric quasi-hermitian models with nontrivial domains of observability. The boundaries of such domains are formed of the so-called exceptional points (EPs) [12, 13], points in parameter space, where the operator ceases to be diagonalizable. The boundary crossings of physical parameters (e.g. time) are studied under the name quantum catastrophes [14], which emphasizes close relationship to classical catastrophe theory [15].

This paper is divided as follows: in section 2, we introduce the general case of the examined $\mathcal{PT}$ -symmetric SSH model. In section 3, we link its special case $\cos \theta = 0$ to a discretized Robin square well, and construct a complete set of pseudometrics. In section 4, we repeat this procedure for a dual SSH nearest-neighbor interaction. In section 5, we consider the general case $H = H_{SSH} + H_I$, and address the questions of pseudometric construction and cutoff emergence. Section 6 is devoted to discussion and remarks.

2 The $\mathcal{PT}$ -symmetric SSH model

Inspired by [2], we examine a SSH model with open boundary conditions, complemented by the $\mathcal{PT}$ -symmetric boundary term as in eq. 2. Throughout this paper, we set with no loss of generality $t = 1$, $\Delta = 1$, and denote $\lambda = \cos \theta$ in eq. 1. The coupling constant γ was taken to be purely imaginary in [2], while we assume a more general case $\gamma = \rho + i\omega \in \mathbb{C}$. In order to apply powerful tools of linear algebra, we work in a matrix representation of the creation and

annihilation operators. The basis of a corresponding two-dimensional Hilbert space is chosen, such that

$$\begin{aligned} a|0\rangle &= 0, & a^\dagger|0\rangle &= |1\rangle \\ a|1\rangle &= |0\rangle, & a^\dagger|1\rangle &= 0 \end{aligned} \quad (6)$$

In such case, the model from eq. 1 becomes a family of $n \times n$ matrices with $n = 2N$. We adopt a strategy of printing the matrices explicitly for the $n = 4$, as long as the extrapolation pattern for higher $n \in \mathbb{N}$ is clear enough. That means we may write

$$H^{(4)} = H_{SSH}^{(4)} + H_I^{(4)} = \begin{bmatrix} \gamma & -1 - \lambda & & \\ -1 - \lambda & & -1 + \lambda & \\ & -1 + \lambda & & -1 - \lambda \\ & & -1 - \lambda & \gamma^* \end{bmatrix} \quad (7)$$

In parallel with numerical experiments of [2], we illustrate the effects of general complex coupling in fig. I. Clearly, there is no reason to confine attention to purely imaginary γ in our $\mathcal{PT}$ -symmetric considerations, as for ρ reasonably small, there always exists a nonempty interval of ω with real spectrum. Size of such interval shrinks with growing ρ , and finally vanishes completely, in the present case for $\rho \approx 1$.




Figure I: $Re[\sigma(H)]$ and $Im[\sigma(H)]$ for the $n = 50$ SSH model as a function of θ . The values of γ are $0.5i, 0.5i + 1$ and $0.5i + 2$.




Figure II: Exceptional points of eq. 8 for $n = 3, 4, 5$ in the (α, β) and (ρ, ω) coordinates.

3 The Robin square well

The construction of Θ for infinite-dimensional quasi-hermitian models is in general a highly nontrivial task, which has to be approached perturbatively. An exceptional operator admitting exact metric construction (Laplacian on a real interval with imaginary Robin boundary conditions) was introduced in [16]. This inspired a subsequent paper [17], which applied discretization techniques on this continuous model, resulting in a $\mathcal{PT}$ -symmetric family of matrices

$$H_{DR}^{(4)} = \begin{bmatrix} \gamma & -1 & & \\ -1 & & -1 & \\ & -1 & & -1 \\ & & -1 & \gamma^* \end{bmatrix} \quad \text{with} \quad \gamma = \frac{1}{1 - \alpha - i\beta} \quad (8)$$

which clearly coincides with eq. 7 for $\lambda = 0$. Throughout this paper, we shall use the natural parametrization $\gamma = \rho + i\omega$ instead of eq. 8. The coordinate transformation connecting these parametrizations is

$$\omega = \frac{\beta}{(1 - \alpha)^2 + \beta^2}, \quad \rho = \frac{1 - \alpha}{(1 - \alpha)^2 + \beta^2} \quad (9)$$

The domains of quasi-hermicity in both coordinate systems are shown in fig. II. The non-shrinking behavior of the domains in the limit $n \rightarrow \infty$ agrees with the existence of a quasi-hermitian operator in the continuous limit. Now, recall that any $n \times n$ metric may be expressed as $\Theta = \sum_{k=1}^n \kappa_n |n\rangle \langle n|$ [18], where $|n\rangle$ are the eigenvectors of $H^\dagger$. Thus, a general metric depends on n arbitrary parameters. Equivalently, we can construct n linearly independent hermitian (but not necessarily positive) matrices $\mathcal{P}^{(k)}$ for a given H , such that

$$\Theta = \sum_{k=1}^n \varepsilon_k \mathcal{P}^{(k)} \quad (10)$$

with ε_k being constrained by the condition $\Theta > 0$. Such a set of $\mathcal{P}^{(k)}$ is called a complete set of pseudometrics. Although the decomposition from eq. 10 may be in general quite arbitrary, our aim is to find the pseudometrics in a form, which admits extrapolation to general $n \in \mathbb{N}$. Already in [17], the authors appreciated the existence of a particular metric Θ with elements

$$\Theta_{jk} = \begin{cases} 1 & j = k \\ -i\omega(\rho - i\omega)^{j-k-1} & j > k \\ i\omega(\rho + i\omega)^{j-k-1} & j < k \end{cases} \quad (11)$$

which served also as a starting point of our considerations. For sufficiently low dimensions, we can construct pseudometrics explicitly with the help of symbolic manipulation on any computer algebra system. For the present two-site model, after denoting $\xi = \rho - i\omega$, we get

$$\begin{aligned}
\mathcal{P}^{(1)} &= \begin{bmatrix} 1 & -i\omega & -i\omega\xi & -i\omega\xi^2 \\ i\omega & 1 & -i\omega & -i\omega\xi \\ i\omega\xi^* & i\omega & 1 & -i\omega \\ i\omega\xi^{*2} & i\omega\xi^* & i\omega & 1 \end{bmatrix} & \mathcal{P}^{(2)} &= \begin{bmatrix} 1 & 1 & -i\omega & -i\omega\xi \\ 1 & \rho & 1 & -i\omega \\ i\omega & 1 & \rho & 1 \\ i\omega\xi^* & i\omega & 1 & \end{bmatrix} \\
\mathcal{P}^{(3)} &= \begin{bmatrix} 1 & -i\omega \\ 1 & \rho & 1 \\ 1 & \rho & 1 \\ i\omega & 1 \end{bmatrix} & \mathcal{P}^{(4)} &= \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}
\end{aligned} \tag{12}$$

Note that $\mathcal{P}^{(1)}$ coincides with [eq. 11](#), while $\mathcal{P}^{(4)}$ realizes a discrete operator of parity, and the condition $H^\dagger \mathcal{P}^{(4)} = \mathcal{P}^{(4)} H$ demonstrates the $\mathcal{PT}$ -symmetry of the present model. The two-site case suggests an extrapolation pattern for higher $n \in \mathbb{N}$ with the k -th pseudometric having only $2(n - k) + 1$ nonzero antidiagonals. The nonzero elements of such metrics are given by the following table

$-i\omega\xi^{(i-j-k)}$	$i - j \geq k$
$i\omega\xi^{*(i-j-k)}$	$j - i \geq k$
ρ	$ i - j < k, i + j - k \text{ even}$
1	$ i - j < k, i + j - k \text{ odd}$

(13)

A rigorous proof of the above proposition by double induction $n \rightarrow n + 1$ and $k \rightarrow k + 1$ would be a straightforward, although lengthy, generalization of proposition 2 in [\[17\]](#) (which is essentially the proof of $n \rightarrow n + 1$ for $k = 1$), and therefore may be omitted. The positivity of the resulting metric as a function of ε_i is a generally nontrivial problem, which lies outside of the scope of the present discussion. We can however make use of the fact, that the pseudometric $\mathcal{P}^{(1)}$ is a genuine metric, and treat other ε_i using tools of perturbation theory.

4 The dual-SSH model

A large subclass of tridiagonal quasi-hermitian matrices has a close connection to the theory of orthogonal polynomials [\[19, 20\]](#). Inspired by these models, we define the dual-SSH (or dSSH) model, which, in parallel with the operator-matrix correspondence of [eq. 7](#) has its two-site form

$$H_{dSSH}^{(4)} = \begin{bmatrix} \gamma & -1 - \lambda & & \\ -1 + \lambda & & -1 - \lambda & \\ & -1 + \lambda & & -1 - \lambda \\ & & -1 + \lambda & \gamma^* \end{bmatrix} \tag{14}$$

Schematic drawings of the both dSSH and the original SSH interaction are shown for $N = 4$ in [fig. III](#), with black circle denoting the boundary term from [eq. 2](#), and solid, respectively dashed lines denoting the $-1 - \lambda$, respectively $-1 + \lambda$ interaction

In this section, we study the special case of [eq. 14](#) with $\gamma = 0$. This matrix family indeed belongs to a class of tridiagonal models admitting elegant treatment using theory of orthogonal polynomials. Recall that any orthogonal polynomial sequence $p_n(x)$ obeys a three-term recurrence relation




Figure III: *Schemes of the four-site SSH and dSSH models.*

$$xp_n(x) = a_{n(n+1)}p_{n+1}(x) + a_{nn}p_n(x) + a_{n(n-1)}p_{n-1}(x) \quad (15)$$

with $a_{n(n+1)}a_{(n+1)n} > 0$ and $a_{nn} \in \mathbb{R}$. The coefficients a_{ij} may be understood as elements of truncated three-diagonal $n \times n$ matrices with characteristic polynomial $p_n(x)$. Consequently, the eigenvalues of such matrices (the roots of p_n) are real and distinct, which is a sufficient condition for matrix quasi-hermicity. Moreover [21], a complete set of pseudometrics for such models has a very feasible form, with $\mathcal{P}^{(k)}$ having only $(2k + 1)$ nonzero diagonals, and may be constructed by solving the recurrences

$$\sum_{j=0}^{k-1} a_{(k+j)(k)} \mathcal{P}_{(k+j)(k+1)}^{(k)} = \sum_{j=0}^{k-1} a_{(k+j)(k+1)} \mathcal{P}_{(k)(k+j)}^{(k)} \quad (16)$$

The matrix elements of eq. 14 obey the condition $a_{n(n+1)}a_{(n+1)n} > 0$ for any $\theta \neq k\pi$. At the exceptional points $\theta = k\pi$ the operator is not diagonalizable despite the reality of its spectrum, and its quantum-mechanical interpretation is lost. For $\theta \neq k\pi$, we may construct pseudometrics using eq. 16, or again with the help of computer-based symbolic manipulations. In either case, we arrive at the two-site formulae

$$\mathcal{P}^{(1)} = \begin{bmatrix} + & & & \\ & - & & \\ & & + & \\ & & & - \end{bmatrix} \quad \mathcal{P}^{(2)} = \begin{bmatrix} & 1 & & \\ 1 & & 1 & \\ & 1 & & 1 \\ & & & 1 \end{bmatrix} \quad \mathcal{P}^{(3)} = \begin{bmatrix} & & + & \\ & - & & - \\ + & & + & \\ & - & & \end{bmatrix} \quad \mathcal{P}^{(4)} = \begin{bmatrix} & & & 1 \\ & & 1 & \\ & 1 & & \\ 1 & & & \end{bmatrix} \quad (17)$$

where we have denoted $\pm = (1 \pm \lambda)$. These formulae suggest an extrapolation pattern with k -th pseudometric having $2k + 1$ nonzero diagonals and $2(n - k) + 1$ nonzero antidiagonals. Moreover, the nonzero elements are arranged in a chessboard-like pattern, in accordance with [20]. We may write formula for zero elements as

$$\mathcal{P}_{ij}^{(k)} = 0 \begin{cases} \text{for } |i + j - k| \text{ odd} \\ \text{for } |i - j| \geq k \text{ or } |i + j - n - 1| > n - k \end{cases} \quad (18)$$

Nonzero elements of $\mathcal{P}^{(k)}$ are given by the following table. In the left column, we have listed for comparison also nonzero pseudometric elements for the classical SSH interaction from eq. 7 with $\gamma = 0$. The hermicity of such model is apparent from $\Theta = I$ being among the admissible metrics.

SSH		dSSH	
+	k even, i odd	+	k odd, i odd
-	k even, i even	-	k odd, i even
1	k odd	1	k even

(19)

5 The inner products

After examining two special cases $\gamma = 0$ and $\lambda = 0$, we are ready to address the model of [eq. 7](#) in full generality. In order to print the resulting matrices explicitly, we define $\pm = (1 \pm \lambda)w$, where w appearing in the (ij) -th element of the k -th pseudometric denotes the (ij) -th element of the k -th pseudometric of [eq. 12](#). Explicit construction of metric operators remains feasible for sufficiently low dimensions. For the two-site model, we get the following results (we shall explain shortly why $\mathcal{P}^{(1)}$ is missing from the list)

$$\begin{aligned}
\mathcal{P}_{SSH}^{(2)} &= \begin{bmatrix} + & - & + & 1 \\ + & - & -^2 & + \\ + & -^2 & - & +- \\ 1 & + & +- & \end{bmatrix} & \mathcal{P}_{SSH}^{(3)} &= \begin{bmatrix} + & 1 \\ - & 1 & + \\ + & 1 & - \\ 1 & + & \end{bmatrix} & \mathcal{P}_{SSH}^{(4)} &= \begin{bmatrix} 1 \\ 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \\
\mathcal{P}_{dSSH}^{(2)} &= \begin{bmatrix} -^2 & - & 1 \\ -^2 & - & +- & + \\ - & +- & + & +^2 \\ 1 & + & +^2 & \end{bmatrix} & \mathcal{P}_{dSSH}^{(3)} &= \begin{bmatrix} - & 1 \\ - & 1 & + \\ - & 1 & + \\ 1 & + & \end{bmatrix} & \mathcal{P}_{dSSH}^{(4)} &= \begin{bmatrix} 1 \\ 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}
\end{aligned}
\tag{20}$$

Extrapolation formulae for general $n \in \mathbb{N}$ are clear from these matrices, again with the k -th pseudometric having $2(n-k)+1$ nonzero antidiagonals. However, as long as we wish to impose the same ansatz for pseudometrics $\mathcal{P}^{(k)}$, $k < n-2$, we discover the emergence of a cutoff value, for which the pattern ceases to be valid. This phenomenon can be seen already in the case of hepta-antidiagonal pseudometrics. We illustrate this behavior on the following three-site models

$$\begin{aligned}
\mathcal{P}_{SSH}^{(3)} &= \begin{bmatrix} +^2 & +- & + & 1 \\ +-^2 & +- & +^2 & -^2 & + \\ +^2 & +- & +-^2 & \Lambda_1 & +^2 & +- \\ +- & +^2 & \Lambda_1 & +-^2 & +- & +^2 \\ + & -^2 & +^2 & +- & +-^2 & \\ 1 & + & +- & +^2 & \end{bmatrix} & \mathcal{P}_{dSSH}^{(3)} &= \begin{bmatrix} -^3 & -^2 & - & 1 \\ -^3 & +- & +^2 & +- & + \\ -^3 & +- & +-^2 & \Lambda_2 & +^2 & +^2 \\ -^2 & +^2 & \Lambda_2 & +-^2 & +- & +^3 \\ - & +- & +^2 & +- & +^3 & \\ 1 & + & +^2 & +^3 & \end{bmatrix}
\end{aligned}
\tag{21}$$

where $+^{-2}$ stands for $+(-)^2$. If we insert these matrices into the equation $H^\dagger \mathcal{P}^{(3)} = \mathcal{P}^{(3)} H$, we obtain complex solutions for Λ_i , which violate the hermicity condition. Thus, in the general case of [eq. 7](#), we are able to build just three pseudometric families $\mathcal{P}^{(n-2)}$, $\mathcal{P}^{(n-1)}$ and $\mathcal{P}^{(n)}$ before cutoff emergence. Still, the pattern of matrix elements not approaching infinity as $n \rightarrow \infty$ is strongly conjectured to be preserved, suggesting that the resulting metrics Θ remain bounded in appropriate continuous limit. Such a continuous analogue is well-known for $\lambda = 0$ [\[16\]](#), but the examination of the continuous limit for general $\theta \neq 0$ remains open.

6 Discussion

We have successfully constructed a complete set of pseudometrics for the discrete Robin square well, as well as for the hermitian SSH model and its dual counterpart. In the general case of [eq. 1](#), we encountered a cutoff preventing the construction of pseudometrics beyond three particular families. While this may be sufficient in many scenarios, we emphasize two reasons for the importance of constructing a complete set of pseudometric. As long as the Hamiltonian is not the only dynamical observable, the presence of additional observables Λ_i imposes additional constraints on the metric, in the form of equations

$$\Lambda_i^\dagger \Theta = \Theta \Lambda_i \quad (22)$$

Furthermore, the physical Hilbert spaces $\mathcal{H}^{(S)}$ generated by different metrics are in general not unitarily equivalent. In particular, for parametric models, some metrics may have larger domains of positivity than others [\[22, 23\]](#). The choice of a correct physical metric is then dictated by other physical principles of, for example, locality.




Figure IV: *Eigenvalues of the $n = 50$ SSH, dSSH and the model $-1 - \lambda \rightarrow -1 + \lambda$ for $\gamma = 0.5i + 0.8$.*

Spectral behavior of the discussed models is briefly outlined for a single parameter value in [fig. IV](#), where we have, for comparison, shown also the operator obtained from [eq. 7](#) by transforming all elements $-1 - \lambda$ into $-1 + \lambda$. All three models exhibit nontrivial domains of observability occurring for non-zero γ . For non-hermitian operators, it may show rewarding, in addition to studying the spectrum, to examine their ε -pseudospectrum [\[24, 25\]](#)

$$\sigma_\varepsilon(H) = \{\lambda \in \mathbb{C} \mid \|(H - \lambda)^{-1}\| \geq \varepsilon^{-1}\} \quad (23)$$

While the pseudospectrum of a general operator may behave quite arbitrarily, pseudospectra of quasi-hermitian operators admit a simple characterization based on the spectral theorem. Using the decomposition $\Theta = \Omega^\dagger \Omega$, we may write

$$\frac{1}{\rho(\lambda, \sigma(H))} \leq \|(H - \lambda)^{-1}\| \leq \frac{\|\Omega\| \|\Omega^{-1}\|}{\rho(\lambda, \sigma(H))} \quad (24)$$

therefore the ε -pseudospectrum is contained in the $\kappa\varepsilon$ -neighborhood of the spectrum, with $\kappa = \|\Omega\| \|\Omega^{-1}\|$. Clearly, the pseudospectrum contains information unavailable from the spectrum itself, with the resolvent norm being pronounced near the eigenvalues, which are likely to complexify by a small perturbation of parameters.




Figure V: Pseudospectra of [eq. 8](#) for $n = 8$ and $\alpha = 1, \beta = 1$, respectively $\alpha = 1, \beta = 0.7$

References

- [1] W. P. Su, J. R. Schrieffer, and A. J. Heeger. “Solitons in polyacetylene”. *Phys. Rev. Lett.* 42 (25 1979), pp. 1698–1701.
- [2] B. Zhu, R. Lu, and S. Chen. “ $\mathcal{PT}$ -symmetry in the non-hermitian Su-Schrieffer-Heeger model with complex boundary potentials”. *Phys. Rev. A* 89 (2014), p. 062102. arXiv:[1405.5591](#) [[cond-mat.other](#)].
- [3] P. Delplace, D. Ullmo, and G. Montambaux. “Zak phase and the existence of edge states in graphene”. *Phys. Rev. B* 84 (2011), p. 195452. arXiv:[1109.4608](#) [[cond-mat](#)].
- [4] Z. Zhang et al. “Modulated pair condensate of p-orbital ultracold fermions”. *Phys. Rev. A* 82 (2010), p. 033610. arXiv:[0910.2431](#) [[cond-mat.quant-gas](#)].
- [5] S. Ryu et al. “Topological insulators and superconductors: tenfold way and dimensional hierarchy”. *New J. Phys.* 12 (2010), p. 065010. arXiv:[0912.2157](#) [[cond-mat](#)].
- [6] C. M. Bender and S. Boettcher. “Real spectra in non-hermitian Hamiltonians having $\mathcal{PT}$ -symmetry”. *Phys. Rev. Letters* 80 (1998), pp. 5243–5246. arXiv:[physics/9712001](#).
- [7] C. M. Bender. “Making sense of non-hermitian Hamiltonians”. *Rep. Prog. Phys.* 70 (2007), pp. 947–1018. arXiv:[hep-th/0703096](#).
- [8] P. Dorey, C. Dunning, and R. Tateo. “The ODE/IM correspondence”. *J. Phys. A: Math. Theor.* 40 (2007), R205. arXiv:[hep-th/0703066](#).
- [9] M. Znojil. “Three-Hilbert-space formulation of quantum mechanics”. *SIGMA* 5 (2009), p. 1. arXiv:[0901.0700](#) [[quant-ph](#)].
- [10] F. G. Scholtz, H. B. Geyer, and F. J. W. Hahne. “Quasi-hermitian operators in quantum mechanics and the variational principle”. *Ann. Phys.* 213 (1992), pp. 74–101.
- [11] A. Mostafazadeh. “Metric operators for quasi-hermitian Hamiltonians and symmetries of equivalent hermitian Hamiltonians”. *Journal of Physics A Mathematical General* 41 (2008), p. 244017. arXiv:[0707.3075](#) [[quant-ph](#)].
- [12] T. Kato. *Perturbation theory for linear operators*. Springer, 1995.
- [13] W. D. Heiss. “The physics of exceptional points”. *J. Phys. A: Math. Theor.* 45 (2012), p. 4016. arXiv:[1210.7536](#) [[quant-ph](#)].
- [14] M. Znojil. “Quantum catastrophes: a case study”. *J. Phys. A: Math. Theor.* 45 (2012), p. 4036. arXiv:[1206.6000](#) [[quant-ph](#)].

- [15] V. Arnold. *Catastrophe Theory*. Springer, 1992.
- [16] D. Krejcirik, H. Bila, and M. Znojil. “Closed formula for the metric in the Hilbert space of a $\mathcal{PT}$ -symmetric model”. *J. Phys. A: Math. Gen.* 39 (2006), pp. 10143–10153. arXiv:[math-ph/0604055](#).
- [17] M. Znojil. “Solvable non-hermitian discrete square well with closed-form physical inner product”. *J. Phys. A: Math. Theor.* 47 (2014), p. 435302. arXiv:[1409.3788v1 \[quant-ph\]](#).
- [18] M. Znojil. “On the role of the normalization factors κ_n and of the pseudometric $P \neq P^\dagger$ in crypto-hermitian quantum models”. *SIGMA* 4 (2008), p. 1. arXiv:[0710.4432 \[math-ph\]](#).
- [19] G. Lévai, F. Ruzicka, and M. Znojil. “Three solvable matrix models of a quantum catastrophe”. *Int. J. Theor. Phys.* 53 (2014), pp. 2875–2890. arXiv:[1403.0723 \[quant-ph\]](#).
- [20] M. Znojil and J. Wu. “A generalized family of discrete $\mathcal{PT}$ -symmetric square wells”. *Int. J. Theor. Phys.* 52 (2013), pp. 2152–2162. arXiv:[1302.1662 \[quant-ph\]](#).
- [21] M. Znojil. “Quantum inner-product metrics via the recurrent solution of the Dieudonné equation”. *J. Phys. A: Math. Theor.* 45 (2012), p. 085302. arXiv:[1201.2263 \[math-ph\]](#).
- [22] M. Znojil and D. Borisov. “Mathematical and physical meaning of the crossings of energy levels in $\mathcal{PT}$ -symmetric systems”. *ArXiv e-prints* (2015). arXiv:[1303.4876 \[quant-ph\]](#).
- [23] D. I. Borisov, F. Ruzicka, and M. Znojil. “Multiply Degenerate Exceptional Points and Quantum Phase Transitions”. *Int. J. Theor. Phys.* 54 (2015), pp. 4293–4305. arXiv:[1412.6634 \[quant-ph\]](#).
- [24] L. N. Trefethen and M. Embree. *Spectra and pseudospectra - the behavior of nonnormal matrices*. Princeton University Press, 2005.
- [25] D. Krejcirik et al. “Pseudospectra in non-hermitian quantum mechanics”. *J. Math. Phys.* 56 (2014), p. 103513. arXiv:[1402.1082 \[math.SP\]](#).